

YOUTH DEVELOPMENT PROGRAM EVALUATION

ANNUAL REPORT

PREPARED BY:

IOWA CONSORTIUM FOR SUBSTANCE ABUSE RESEARCH AND EVALUATION UNIVERSITY OF IOWA, IOWA CITY, IOWA 52242-5000

WITH FUNDS PROVIDED BY:

U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES, SUBSTANCE ABUSE AND MENTAL HEALTH SERVICES ADMINISTRATION, CENTER FOR SUBSTANCE ABUSE PREVENTION, IOWA DEPARTMENT OF PUBLIC HEALTH, BUREAU OF SUBSTANCE ABUSE PREVENTION AND TREATMENT

CITATION OF REFERENCES RELATED TO THIS REPORT IS APPRECIATED. SUGGESTED CITATION:

WHITE, K., CLAYTON, R., & ARNDT, S. (2009). YOUTH DEVELOPMENT PROGRAM EVALUATION: ANNUAL REPORT. (IOWA DEPARTMENT OF PUBLIC HEALTH CONTRACT #5889YM50). IOWA CITY, IA: IOWA CONSORTIUM FOR SUBSTANCE ABUSE RESEARCH AND EVALUATION. <u>http://iconsortium.subst-abuse.uiowa.edu/</u>



YOUTH DEVELOPMENT PROGRAM EVALUATION

ANNUAL REPORT

JULY 2008 - JUNE 2009

PREPARED BY:

KRISTIN WHITE, MA PROGRAM EVALUATOR

REBECCA CLAYTON, BS ASSOCIATE DIRECTOR

STEPHAN ARNDT, PHD DIRECTOR

© 2009 UNIVERSITY OF IOWA

Table of Contents

Intro	duction .		1
Dem	ographic	S	1
Outc	ome Dat	a	2
Past	30-Day	Use	3
Т	able 1:	Change in Past 30-Day Use: Youth Development Total and Results by Program	3
G	Graph 1:	Change in Past 30-Day Use: Youth Development Total and Results by Program	4
Attitu	ides Tow	ard Substance Use	5
G	Graph 2:	Change in Attitudes Toward Substance Use: Youth Development	5
G	Graph 3:	Change in Attitudes Toward Substance Use: LifeSkills Training	6
G	Graph 4:	Change in Attitudes Toward Substance Use: Project Alert	6
Perc	eived Ris	sk of Harm from Substance Use	7
G	Graph 5:	Change in Perceived Risk of Harm: Youth Development	7
G	Graph 6:	Change in Perceived Risk of Harm: LifeSkills Training	8
G	Graph 7:	Change in Perceived Risk of Harm: Project Alert	8
Conc	clusion		9
т	able 2:	Positive Outcome Percentages for Attitudes Toward Substance Use	9
Т	able 3:	Positive Outcome Percentages for Perceived Risk of Harm from Use	10

Introduction

The purpose of the Youth Development project is to provide evidence-based substance abuse prevention programming for youth ages 5 through 18 that includes in- and out-of-school opportunities for youth development, character development and youth leadership. Nine substance abuse prevention organizations participate in this project: Clinton Substance Abuse Council; Community and Family Resources; Henry County Extension; Jackson Recovery Centers; Loess Hills Area Education Agency; Mason City Youth Task Force; Substance Abuse Treatment Unit of Central Iowa; United Action for Youth; and Youth and Shelter Services.

Evaluation Design

The evaluation employed a matched pre-post design whereby a survey was administered at the beginning and at the conclusion of the prevention program. The post-test survey also is administered after each year of programming for programs spanning multiple years. Post-test survey data from the previous program year is used as a baseline for reporting current year outcomes for multi-year programs. A total of 770 pre-tests and 1475 first, second, and third year post-tests were collected for the Youth Development project in State Fiscal Year 2009. This resulted in a total of 915 survey matches for analysis. Of those, 505 were from participants in single-year programs and participants in the first year of multi-year programs; 352 were from participants in the second year of multi-year programs; and 58 were from participants in the third year of multi-year programs.

The pre-post data were used to help answer the following evaluation questions:

- Has alcohol/tobacco/marijuana usage changed in the target population?
- Has the percentage of the target population who indicate positive attitudes at baseline (pre-test) maintained or increased after the intervention (post-test)?
- Has perceived risk of harm from alcohol/tobacco/marijuana use maintained a positive response or increased from pre-test to post-test?

Demographics

The median age of Youth Development project participants at post-test was twelve years. More than three-quarters of participants (76.5%) were in the sixth and seventh grades. The sex of participants was split almost equally, with 50.4% being female. Nearly one quarter (23.9%) of participants were Hispanic or Latino. Participant racial groups are delineated below:

- 78.5% White;
- 2.7% Black/African American;
- 2.0% American Indian/Alaska Native;
- 1.0% Asian;
- 0.6% Hawaiian/Other Pacific Islander;
- 0.1% Arab American; and
- 15.2% more than one race.

Outcome Data

The figures included in the outcome data section detail past 30-day use, attitudes toward use, and perceptions of risk of harm from use of alcohol, tobacco, and marijuana. (Please note: the past 30-day use questions for tobacco include cigarettes, snuff, and cigars, whereas the attitude and perceived risk questions include only cigarettes.) Figures are provided for the project total (all participants in the Youth Development project for whom matched surveys existed in the data set), for participants in the LifeSkills Training (LST) program, and for participants in Project Alert. Other programs did not meet the minimum sample size necessary for individual reporting, but are included in the Youth Development project totals. Those programs are: Leadership and Resiliency Program; Reconnecting Youth; Teen Task Force; Teen Club; Service Learning Initiative Project (SLIP); and Positive Action.

Due to the number of unmatched pre-tests (29%, possibly caused by survey administration anomalies, data entry error, early termination of programming, or ID assignment mistakes) an attrition analysis was performed on these data. Grade and attitude toward alcohol use had statistically significant associations with whether an individual completed a post-test (Cochran-Mantel-Haenszel tests, p<0.05). Participants in the seventh grade tended to drop out of programming or not complete a post-test. There also was a slight tendency for participants who thought alcohol use was only a little wrong or not wrong at all to drop out or not complete a post-test. Therefore, findings may apply only to those youth who completed the program and completed a post-test, rather than all participants.

lowa Youth Survey (IYS) data are provided as a reference point for interpreting the outcome data in this report. The Iowa Youth Survey is a triennial census assessment of Iowa's secondary school-age students' (grades 6, 8, and 11) attitudes toward substance use and actual use of substances. The 2008 IYS data included here provide an estimate of the change one might expect to see each year in Iowa's general youth population due to maturation. Thus, IYS data serve as a general point of reference when examining Youth Development program outcomes (change from pre-test to post-test) rather than comparing program percentages to zero, or no change. (Note: Youth who participated in Youth Development programming may also have completed the IYS).

Past 30-Day Use

Table 1, below, and Graph 1, on page 4, present data on past 30-day use of alcohol, tobacco, and marijuana for all Youth Development participants (YD Total), for LifeSkills Training participants and for Project Alert participants. A positive (+) percent change indicates an increase in use; whereas a negative (-) change indicates a decrease in use. Past 30-day use of marijuana decreased from pre-test to post-test in project participants overall. Past 30-day use of alcohol and marijuana also decreased among Project Alert participants. Past 30-day use of tobacco increased for all groups, and past 30-day use of all three substances increased for LifeSkills Training participants. However, all increase percentages are lower than the IYS percentages.

Group	N	Median Age	Alcohol		Tobacco		Marijuana	
			Pre-Test %	Change	Pre-Test %	Change	Pre-Test %	Change
IYS ¹	64,937 ²	13	_	+5.00	_	+2.50	_	+1.00
YD Total	915	12	4.09	+0.22	2.65	+0.78	1.55	-0.22
LST	661	12	1.87	+0.47	0.31	+0.94	0	+0.62
PA	173	12	2.96	-0.59	0.59	+1.78	0.59	-0.59


Percentage of Youth Reporting Past 30-Day Use at the Pre-Test and Change at Post-Test

Table 1. Change in Past 30-Day Use: Youth Development Total and Results by Program

¹ IYS entries indicate the yearly average change in 30-day use between all lowa students in grades 6 and 8. The majority of 6th graders completing the IYS were 11 years old; the majority of 8th graders were 13. Data were from the 2008 lowa Youth Survey, State of lowa report (pgs.12, 29-30, and 90-91).

²The total number of 6th graders completing the lowa Youth Survey was 32,264; the total number of 8th graders was 32,673.

Key				
IYS	Iowa Youth Survey	LST	LifeSkills Training	
YD Total	All Youth Development Participants	PA	Project Alert	






<u>Key</u>				
IYS	Iowa Youth Survey	LST	LifeSkills Training	
YD Total	All Youth Development Participants	PA	Project Alert	


Attitudes Toward Substance Use

Graphs 2 through 4 on pages 5 and 6 show change in individual attitudes from the pre-test to the post-test for alcohol, cigarettes, and marijuana. Individual attitudes either: 1) "improved," which means that attitudes grew more unfavorable toward use of alcohol, tobacco, or marijuana (e.g., respondent felt alcohol use was wrong at pretest and very wrong at post-test); 2) "maintained +," which means that the pre- and post-test responses remained the same and were unfavorable toward alcohol. tobacco, or marijuana use (a positive outcome); 3) "maintained -," which means that the pre- and post-test responses remained the same and were favorable toward alcohol, tobacco, or marijuana use (a negative outcome); or 4) "worsened," meaning that attitudes grew more favorable toward alcohol, tobacco, or marijuana use from pretest to post-test (e.g., respondent felt marijuana use was very wrong at pre-test and a little bit wrong at post-test). Desired outcomes for these questions are improvement in ("improved") or positive maintenance ("maintained +") of attitudes toward substance use. Graphs 2, 3, and 4 show data for all Youth Development project participants, LifeSkills Training participants, and Project Alert participants, respectively. The number of matched pre- and post-tests containing a response to the question is shown in the key, following each substance.



Graph 2. Change in Attitudes Toward Substance Use: Youth Development


Youth Development project participants showed positive outcomes as follows: Alcohol – 87%; Cigarettes – 93%; and Marijuana – 90%.



Graph 3. Change in Attitudes Toward Substance Use: LifeSkills Training

LifeSkills Training participants showed positive outcomes as follows: Alcohol – 88%; Cigarettes – 94%; and Marijuana – 90%.


Graph 4. Change in Attitudes Toward Substance Use: Project Alert



Project Alert participants showed positive outcomes as follows: Alcohol – 90%; Cigarettes – 96%; and Marijuana – 94%.


Perceived Risk of Harm from Substance Use

Graphs 5 through 7 show change from pre- to post-test in individuals' perceptions of risk of harm from use of alcohol, cigarettes, and marijuana. Perceptions of risk either: 1) "increased." which means that their reported perceived risk regarding alcohol. tobacco, or marijuana use increased from pre-test to post-test (e.g., respondent felt alcohol use was a moderate risk at pre-test and a great risk at post-test); 2) "maintained +," which means that the pre- and post-test responses remained the same and were unfavorable toward alcohol, tobacco, or marijuana use (a positive outcome); 3) "maintained -," which means that the pre- and post-test responses remained the same and were favorable toward alcohol, tobacco, or marijuana use (a negative outcome); or 4) "decreased," meaning that their reported perception of risk of harm decreased from pre-test to post-test (e.g., respondent reported that marijuana use posed a moderate risk of harm at pre-test and no risk at post-test). Desired outcomes for these questions are an increase in or positive maintenance ("maintained +") of change in perceived risk. Graphs 5, 6, and 7 show data for all Youth Development project participants, LifeSkills Training participants, and Project Alert participants, respectively. The number of matched pre- and post-tests containing a response to the question is shown in the key, following each substance.



Graph 5. Change in Perceived Risk of Harm: Youth Development


Youth Development project participants showed positive outcomes as follows: Alcohol – 82%; Cigarettes – 86%; and Marijuana – 90%.



Graph 6. Change in Perceived Risk of Harm: LifeSkills Training

LifeSkills Training participants showed positive outcomes as follows: Alcohol – 82%; Cigarettes – 86%; and Marijuana – 90%.

Graph 7. Change in Perceived Risk of Harm: Project Alert



Project Alert participants showed positive outcomes as follows: Alcohol – 85%; Cigarettes – 91%; and Marijuana – 96%.

Conclusion

This evaluation of the Youth Development project answers the following questions:

Has alcohol/tobacco/marijuana usage changed in the target population?

Past 30-day use of marijuana decreased from pre-test to post-test in Youth Development participants overall, and past 30-day use of alcohol and marijuana also decreased among Project Alert participants. Past 30-day use of tobacco increased for all groups, and past 30-day use of all three substances increased for LifeSkills Training participants. However, all increase percentages are lower than the IYS percentages, suggesting perhaps some benefit of the programs in deflecting the increases normally seen in this age group.

Has the percentage of the target population who indicate positive attitudes at baseline (pre-test) maintained or increased after the intervention (post-test)?

Table 2 presents positive outcome percentages for attitudes toward alcohol, tobacco, and marijuana use for all Youth Development participants, LifeSkills Training Participants, and Project Alert Participants. At least 87% of participants in all groups maintained or increased positive attitudes regarding substance use (i.e., that alcohol, cigarette, and marijuana use is wrong or very wrong) from pre-test to post-test. At least 90% of Project Alert participants maintained or increased positive attitudes regarding use of all three substances. More than 93% of Project Alert participants maintained or increased positive attitudes regarding use of tobacco, and more than 94% maintained or increased positive attitudes regarding use of marijuana. In all groups, participants felt that tobacco use was more wrong than alcohol or marijuana use.

Positive Outcome Percentages for Attitudes Toward Substance Use						
Participant Group	Alcohol	Tobacco	Marijuana			
Youth Development Participants	87%	93%	90%			
LifeSkills Training Participants	88%	94%	90%			
Project Alert Participants	90%	96%	94%			

Table 2. Positive Outcome Percentages for Attitudes Toward Substance Use by Participant Group

Has perceived risk of harm from alcohol/tobacco/marijuana use maintained a positive response or increased from pre-test to post-test?

Table 3 presents positive outcome percentages for perceived risk of harm from use of alcohol, tobacco, and marijuana for all Youth Development participants, LifeSkills Training Participants, and Project Alert Participants. At least 82% of participants in all groups maintained or increased positive responses regarding perception of risk of harm from substance use (i.e., that using alcohol, cigarette, or marijuana posed moderate to great risk of harm). Eighty-five percent of Project Alert participants maintained or increased positive responses for all three substances. Ninety-six percent or more of Project Alert participants maintained or increased positive responses for all three substances. Ninety-six percent or more of Project Alert participants maintained or increased positive responses for all three substances. In all groups, participants felt that alcohol use was less risky than tobacco or marijuana use.

Positive Outcome Percentages for Perceived Risk of Harm from Substance Use						
Participant Group	Alcohol	Tobacco	Marijuana			
Youth Development Participants	82%	86%	90%			
LifeSkills Training Participants	82%	86%	90%			
Project Alert Participants	85%	91%	96%			

 Table 3. Positive Outcome Percentages for Perceived Risk of Harm from

 Substance Use by Participant Group